

Biochimica et Biophysica Acta (BBA) -
Biomembranes

Volume 1860, Issue 2, February 2018, Pages 264–271



Study of the mechanism of permeabilization of lecithin liposomes and rat liver mitochondria by the antimicrobial drug triclosan

Konstantin N. Belosludtsev^{a, b, *}, Natalia V. Belosludtseva^a, Kirill S. Tenkov^b, Nikita V. Penkov^c, Alexey V. Agafonov^a, Lyubov L. Pavlik^a, Valery A. Yashin^c, Victor N. Samartsev^b, Mikhail V. Dubinin^b

[Show more](#)

<https://doi.org/10.1016/j.bbamem.2017.09.018>

[Get rights and content](#)

Highlights

- Triclosan is able to permeabilize the membrane of lecithin liposomes.
- Raising pH of the medium inhibits the triclosan-induced liposome permeabilization
- Laurdan fluorescence data indicate a phase heterogeneity in triclosan/lecithin system
- Triclosan promotes a cyclosporin A-insensitive swelling of rat liver mitochondria.
- Triclosan induces cytochrome *c* release from rat liver mitochondria

Abstract

The effect of the antimicrobial compound triclosan (5-chloro-2'-(2,4-dichlorophenoxy)phenol) on the permeability of lecithin liposomes and rat liver mitochondria was studied. It was found that triclosan was able to increase nonspecific permeability of liposomes in a dose-dependent manner, which was detected by the release of the fluorescent probe sulforhodamine B (SRB) from vesicles. A partial release of SRB occurs instantly at the moment of triclosan addition, which is followed by a slow leakage of the dye. The triclosan-induced release of SRB from liposomes grew as pH of the medium was decreased from 9.5 to 7.5. As revealed by the laurdan generalized polarization (GP) technique, triclosan increased laurdan GP in lecithin liposomes, indicating a decrease in membrane fluidity. Measurements of GP as a function of fluorescence excitation wavelength gave an ascending line for triclosan-containing liposomes, which can be interpreted as phase heterogeneity of the lipid/triclosan system. Dynamic light scattering experiments also showed that at a high triclosan-to-lipid molar ratio (~0.5), a population of smaller light-scattering particles (~0.4 of the size of liposomes) appear in the system. Experiments with rat liver mitochondria demonstrated that triclosan (10–70 μM) induced a high-amplitude cyclosporin A-insensitive swelling of the organelles accompanied the release of cytochrome *c*. On the basis of the results obtained, possible mechanisms of the toxic effect of triclosan in eukaryotic cells are discussed.

Article outline

☐ Show full outline

Highlights

Abstract

Graphical abstract

Abbreviations

Keywords

1. Introduction

2. Materials and methods

3. Results

4. Discussion

Transparency document

Acknowledgements

References

Figures and tables








Table 1



Abbreviations

TCS, triclosan; CsA, cyclosporin A; LUV, large unilamellar vesicles; MPT, mitochondrial permeability transition; SRB, sulforhodamine B

Keywords

Triclosan; Membrane permeabilization; Liposomes; Mitochondria

Choose an option to locate/access this article:

Check if you have access through your login credentials or your institution

Purchase \$39.95

Get Full Text Elsewhere

Check Access

Corresponding author at: Institute of Theoretical and Experimental Biophysics, Russian Academy of Sciences, Institutskaya 3, Pushchino, Moscow region 142290, Russia.
© 2017 Elsevier B.V. All rights reserved.

articles

on acyl chain length of energy ...
Chemistry more

ec0005
Effect of surface-potential modulators on the...
2015, Biochimica et Biophysica Acta (BBA) - Biom... more
ADVERTISEMENT

Mer Antimicrobial ...
2008

View

Citin

Rela

Search ScienceDirect

[Advanced search](#)

[About ScienceDirect](#)[Remote access](#)[Shopping cart](#)[Contact and support](#)[Terms and conditions](#)[Privacy policy](#)

Cookies are used by this site. For more information, visit the [cookies page](#).

Copyright © 2017 Elsevier B.V. or its licensors or contributors. ScienceDirect ® is a registered trademark of Elsevier B.V.